

## 7. REFERÊNCIAS BIBLIOGRÁFICAS

1. Cecil Medicina 23 ed (Elsevier). Rio de Janeiro: Elsevier; 2009. 1988-2013.
2. Ministério da Saúde (Brasil), Secretaria de Atenção à Saúde. Consulta Pública Nº 39, de 28 de Outubro de 2010. Brasília: Ministério da Saúde, 2010.
3. Frosch M, Anthony D, De Girolami U. Bases patológicas das doenças; Robbins & Cotran; 7ª Edição; Editora Elsevier, Capítulo 28: O sistema nervoso central, página 1425; 2004.
4. National Clinical Guidelines for Stroke; 2ª edição, Clinical Effectiveness & Evaluation Unit; Royal College of Physicians, 2004; Artigo online - <http://www.sorcan.ca/0C4FC187-741F-412A-81DB-BA1E7920A91F/FinalDownload/DownloadId-2589687BEC0C3AC3D758B8C7D78F17BA/0C4FC187-741F-412A-81DB-BA1E7920A91F/Resources/General/NCG.pdf> (disponível em 05 de Maio de 2011).
5. Inagawa T. Risk factors for the formation and rupture of intracranial saccular aneurysms in Shimane, Japan. *World Neurosurg* 2010;73:155-64.
6. Brisman JL, Song JK, Newell DW. Cerebral aneurysms. *N Engl J Med* 2006; 355: 928-39.
7. Pereira VM, Brina O, Gonzalez AM, Narata AP, Ouared R, Karl-Olof L. Biology and hemodynamics of aneurismal vasculopathies. *Eur J Radiol*. 2013 Jan 21.
8. Gu YX, Chen XC, Song DL, Leng B, Zhao F. Risk factors for intracranial aneurysm in a Chinese ethnic population. *Chin Med J (Engl)* 2006;119:1359-64.
9. Diringner MN. Management of aneurysmal subarachnoid hemorrhage. *Crit Care Med* 2009;37:432-40.

10. De Bock S, Iannaccone F, De Santis G, De Beule M, Mortier P, Verhegghe B, et al. Our capricious vessels: The influence of stent design and vessel geometry on the mechanics of intracranial aneurysm stent deployment. *J Biomech* 2012;45:1353-59.
11. Chung C, Caplan LR. Neurovascular disorders. In: Goetz CG, Pappert EJ (eds). *Textbook of Clinical Neurology*. Pennsylvania, WB Saunders Company, 1999: 907-32.
12. Ojini FI. Natural history of cerebral saccular aneurysms. *West Afr J Med* 2005;24:70-4.
13. Grobelny TJ. Brain aneurysms: epidemiology, treatment options, and milestones of endovascular treatment evolution. *Dis Mon* 2011;57:647–55.
14. Sekhar LN, Heros RC. Origin, growth, and rupture of saccular aneurysms: A review. *Neurosurg* 1981; 8: 248–60.
15. Crawford PM, West CR, Chadwick DW, Shaw MD. Arteriovenous malformations of the brain: natural history in unoperated patients. *J Neurol Neurosurg Psychiatry* 1986;49:1–10.
16. Inci S, Spetzler RF. Intracranial aneurysms and arterial hypertension: A review and hypothesis. *Surg Neurol* 2000; 53: 530–40.
17. Yong-Zhong G, van Alphen HA. Pathogenesis and histopathology of saccular aneurysms: Review of the literature. *Neurol Res* 1990; 12: 249–55.
18. Scanlon A. Aneurysmal subarachnoid haemorrhage – part 1: Pre-operative nursing management diagnosis, complications and treatment: A composite case study. *Contemp Nurse* 2004; 17: 80–8.

19. Shoja MM, Agutter PS, Tubbs RS, Payner TD, Ghabili K, Cohen-Gadol AA. The role of the renin--angiotensin system in the pathogenesis of intracranial aneurysms. *J Renin Angiotensin Aldosterone Syst* 2011;12:262-73.
20. Crawford T. Some observations on the pathogenesis and natural history of intracranial aneurysms. *J Neurol Neurosurg Psychiatry* 1959;22:259-66.
21. Laaksamo E, Ramachandran M, Frösen J, Tulamo R, Baumann M, Friedlander RM, et al. Intracellular signaling pathways and size, shape, and rupture history of human intracranial aneurysms. *Neurosurgery* 2012;70:1565-72.
22. Selman WR, Tarr RW and Ratcheson RA. Intracranial aneurysms and subarachnoid hemorrhage. In: Bradley WG, Daroff RB, Fenichel GM (eds) *Neurology in Clinical Practice. 3d ed.* Boston: Butterworth-Heinemann, 2000, pp.1188–1190.
23. Caranci F, Briganti F, Cirillo L, Leonardi M, Muto M. Epidemiology and genetics of intracranial aneurysms. *Eur J Radiol.* 2013 Feb 8. pii: S0720-048X(13)00011-9.
24. Connolly ES, Solomon RA. Management of unruptured aneurysms. In: Le Roux, P.D., Winn, H.R., Newell, D.W. (Eds.), *Management of Cerebral Aneurysms.* Saunders, Philadelphia, pp. 271–285, 2004.
25. Inagawa T, Hirano A. Autopsy study of unruptured incidental intracranial aneurysms. *Surg Neurol* 1990;34:361–65.
26. Ohaegbulam SC, Dujovny M, Ausman JI, Diaz FG, Malik GM. Ethnic distribution of intracranial aneurysms. *Acta Neurochir (Wien)* 1990;106:132-35.
27. Winn HR, Jane JA Sr, Taylor J, Kaiser D, Britz GW. Prevalence of asymptomatic incidental aneurysms: Review of 4568 arteriograms. *J Neurosurg* 2002; 96:43-49.

28. International Study of Unruptured Intracranial Aneurysms Investigators (ISUIA). Unruptured intracranial aneurysms – risk of rupture and risks of surgical, intervention. *N England J Med* 1998;339:1725–33.
29. Linn FH, Rinkel GJ, Algra A, van Gijn J. Incidence of subarachnoid hemorrhage: role of region, year, and rate of computed tomography: a meta-analysis. *Stroke* 1996; 27:625-29.
30. Inagawa T, Ishikawa S, Aoki H, Takahashi M, Yoshimoto H. Aneurysmal subarachnoid hemorrhage in Izumo City and Shimane Prefecture of Japan. Incidence. *Stroke* 1988;19:170-5.
31. Inagawa T, Tokuda Y, Ohbayashi N, Takaya M, Moritake K. Study of aneurysmal subarachnoid hemorrhage in Izumo City, Japan. *Stroke* 1995;26:761-66.
32. Inagawa T. What are the actual incidence and mortality rates of subarachnoid hemorrhage? *Surg Neurol* 1997;47:47-53.
33. Inagawa T. Trends in incidence and case fatality rates of aneurysmal subarachnoid hemorrhage in Izumo City, Japan, between 1980-1989 and 1990-1998. *Stroke* 2001;32:1499-507.
34. Inagawa T. Seasonal variation in the incidence of aneurysmal subarachnoid hemorrhage in hospital- and community-based studies. *J Neurosurg* 2002;96:497-509.
35. Inagawa T, Takechi A, Yahara K, Saito J, Moritake K, Kobayashi S, et al. Primary intracerebral and aneurysmal subarachnoid hemorrhage in Izumo City, Japan. Part I: incidence and seasonal and diurnal variations. *J Neurosurg* 2000a;93:958-66.

36. Inagawa T, Shibukawa M, Inokuchi F, Tokuda Y, Okada Y, Okada K. Primary intracerebral and aneurysmal subarachnoid hemorrhage in Izumo City, Japan. Part II: management and surgical outcome. *J Neurosurg* 2000b;93:967-75.
37. Inagawa T. Trends in surgical and management outcomes in patients with aneurysmal subarachnoid hemorrhage in Izumo City, Japan, between 1980-1989 and 1990-1998. *Cerebrovasc Dis* 2005;19:39-48.
38. Wiebers DO, Torner JC, Meissner I. Impact of unruptured intracranial aneurysms on public health in the United States. *Stroke* 1992; 23: 1416-19.
39. Xu J, Turner A, Little J, Bleecker ER, Meyers DA. Positive results in association studies are associated with departure from Hardy-Weinberg equilibrium: hint for genotyping error? *Hum Genet* 2002;111:573-4.
42. Inci S, Ozgen T. Multiple aneurysms of the anterior communicating artery. radiological and surgical difficulties. *J Neurosurg* 2005;102:495-502.
40. Phillips LH 2nd, Whisnant JP, O'Fallon WM, Sundt TM Jr. The unchanging pattern of subarachnoid hemorrhage in a community. *Neurology* 1980;30:1034-40.
43. Nguyen TV, Chandrashekar K, Qin Z, Parent AD, Zhang J. Epidemiology of Intracranial Aneurysms of Mississippi: a 10-year (1997-2007) Retrospective Study. *J Stroke Cerebrovasc Dis* 2009; 18:374-80.
44. Amar AP. Brain and Vascular Imaging of Acute Stroke. *World Neurosurg* 2011; 76: S3-S8.
41. Broderick JP, Sauerbeck LR, Foroud T, Huston J 3rd, Pankratz N, Meissner I, et al. The Familial Intracranial Aneurysm (FIA) study protocol. *BMC Med Genet* 2005; 6: 17.

45. Tajiri N, Lau T, Glover LE, Shinozuka K, Kaneko Y, van Loveren H, Borlongan CV. Cerebral aneurysm as an exacerbating factor in stroke pathology and a therapeutic target for neuroprotection. *Curr Pharm Des* 2012;18:3663-69.
46. Keedy A. An overview of intracranial aneurysms. *Mcgill J Med* 2006;9:141–46.
47. Schonfeld AR and McMullen MA. Treatment of brain aneurysms. *Radiol Technol* 2008; 79: 515–31.
48. Burns JD, Brown RD Jr. Treatment of unruptured intracranial aneurysms: surgery, coiling, or nothing? *Curr Neurol Neurosci Rep* 2009; 9: 6–12.
49. Molyneux A, Kerr R, Stratton I, Sandercock P, Clarke M, Shrimpton J, et al. International Subarachnoid Aneurysm Trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised trial. *Lancet* 2002;360:1267–74.
50. Molyneux AJ, Kerr RS, Yu LM, Clarke M, Sneade M, Yarnold JA, International Subarachnoid Aneurysm Trial (ISAT) Collaborative Group. International subarachnoid aneurysm trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised comparison of effects on survival, dependency, seizures, rebleeding, subgroups, and aneurysm occlusion. *Lancet* 2005;366:809–17.
51. Takenaka K, Sakai H, Yamakawa H, Yoshimura S, Kumagai M, Yamakawa H, et al. Polymorphism of the endoglin gene in patients with intracranial saccular aneurysms. *J Neurosurg* 1999; 90: 935-38.
52. Krex D, Schackert HK, Schackert G. Genesis of cerebral aneurysms – an update. *Acta Neurochir (Wien)* 2001; 143: 429-48.

53. Krischek B, Inoue I. The genetics of intracranial aneurysms. *J Human Genet* 2006;51:587–94.
54. Nahed BV, Bydon M, Ozturk AK, Bilguvar K, Bayrakli F, Gunel M. Genetics of intracranial aneurysms. *Neurosurg* 2007;60:213–26.
55. Bilguvar K, Yasuno K, Niemelä M, Ruigrok YM, von Und Zu Fraunberg M, van Duijn CM, et al. Susceptibility loci for intracranial aneurysm in European and Japanese populations. *Nat Genet* 2008;40:1472-77.
56. Yasuno K, Bilguvar K, Bijlenga P, Low SK, Krischek B, Auburger G, et al. Genome-wide association study of intracranial aneurysm identifies three new risk loci. *Nat Genet* 2010; 42: 420–25.
57. Ronkainen A, Hernesniemi J, Puranen M, Niemitukia L, Vanninen R, Ryyanen M, et al. Familial intracranial aneurysms. *Lancet* 1997;349:380–4.
58. Ozüm U, Bolat N, Gül E, Ozdemir O. Endothelial nitric oxide synthase gene [G894T] polymorphism as a possible risk factor in aneurysmal subarachnoid haemorrhage. *Acta Neurochir (Wien)* 2008; 150: 57-61.
59. Schievink WI, Michels VV, Piepgras DG. Neurovascular manifestations of heritable connective tissue disorders. A review. *Stroke* 1994a;25:889-903.
60. Schievink WI, Schaid DJ, Rogers HM, Piepgras DG, Michels VV. On the inheritance of intracranial aneurysms. *Stroke*. 1994b; 25: 2028-37.
61. Schievink WI, Schaid DJ, Michels VV, Piepgras DG. Familial aneurysmal subarachnoid hemorrhage: A community-based study. *J Neurosurg* 1995a;83:426-29.

62. Ronkainen A, Miettinen H, Karkola K, Papinaho S, Vanninen R, Puranen M, et al. Risk of harboring an unruptured intracranial aneurysm. *Stroke* 1998;29:359-62.
63. Chalouhi N, Chitale R, Jabbour P, Tjoumakaris S, Dumont AS, Rosenwasser R, et al. The case for family screening for intracranial aneurysms. *Neurosurg Focus* 2011;31:E8.
64. Goksu E, Akyuz M, Tuncer R. The results of radiological screening in asymptomatic at risk members of intracranial aneurysm families from the Turkish population. *Turk Neurosurg* 2012; 22:55-61.
65. Nahed BV, DiLuna ML, Morgan T, Ocal E, Hawkins AA, Ozduman K, et al. Hypertension, age, and location predict rupture of small intracranial aneurysms. *Neurosurgery* 2005;57:676-83.
66. Ozturk AK, Nahed BV, Bydon M, Bilguvar K, Goksu E, Bademci G, et al. Molecular genetic analysis of two large kindreds with intracranial aneurysms demonstrates linkage to 11q24-25 and 14q23-31. *Stroke* 2006;37:1021-27.
67. Weinsheimer S, Lenk GM, van der Voet M, Land S, Ronkainen A, Alafuzoff I, et al. Integration of expression profiles and genetic mapping data to identify candidate genes in intracranial aneurysm. *Physiol Genomics* 2007;32:45-57.
68. Onda H, Yoneyama T, Akagawa H, Kasuya H. Genetic dissection of intracranial aneurysm. *Brain Nerve* 2008;60:1245-60.
69. Stehbens WE. Familial intracranial aneurysms: an autopsy study. *Neurosurgery* 1998;43:1258-59.
70. Ronkainen A, Niskanen M, Piironen R, Hernesniemi J. Familial subarachnoid hemorrhage. Outcome study. *Stroke* 1999;30:1099-102.



71. Li L, Sima X, Bai P, Zhang L, Sun H, Liang W, et al. Interactions of miR-34b/c and TP53 polymorphisms on the risk of intracranial aneurysm. *Clin Dev Immunol.* 2012;2012:567586.
72. Coutard M, Huang W and Osborne-Pellegrin M. Heritability of intracerebral hemorrhagic lesions and cerebral aneurysms in the rat. *Stroke* 2000; 31: 2678–84.
73. Keramatipour M, McConnell RS, Kirkpatrick P, Tebbs S, Furlong RA, Rubinsztein DC. The ACE I allele is associated with increased risk for ruptured intracranial aneurysms. *J Med Genet* 2000; 37: 498–500.
74. Peck G, Smeeth L, Whittaker J, Casas JP, Hingorani A, Sharma P. The genetics of primary haemorrhagic stroke, subarachnoid haemorrhage and ruptured intracranial aneurysms in adults. *PLoS One* 2008; 3: e3691.
75. Ullrich, DP. Familial cerebral aneurysms including one extracranial internal carotid aneurysm. *Neurology.* 1960 Mar;10:288-94.
76. Graf CJ. Familial intracranial aneurysms. *J Neurosurg* 1966; 25: 304-08.
77. Beumont PJ. The familial occurrence of berry aneurysm. *J Neurol Neurosurg Psychiatry* 1968;31:399-402.
78. Brisman R, Abbassioun K. Familial intracranial aneurysms. *J Neurosurg* 1971; 34:678-82.
79. Edelsohn L, Caplan L, Rosenbaum AE. Familial aneurysms and infundibular widening. *Neurology* 1972; 22: 1056-60.
80. Fox JL, Ko JP. Familial intracranial aneurysms. Six cases among 13 siblings. *J Neurosurg* 1980; 52:501-03.
81. Ronkainen A, Hernesniemi J, Ryyanen M. Familial subarachnoid hemorrhage in east Finland, 1977-1990. *Neurosurgery* 1993;33:787-97.

82. Kojima M, Nagasawa S, Lee YE, Takeichi Y, Tsuda E, Mabuchi N. Asymptomatic familial cerebral aneurysms. *Neurosurg* 1998; 43:776-81.
83. Leblanc R. The risk of intracranial aneurysms in families with subarachnoid hemorrhage. *Can J Neurol Sci* 1995; 22: 333.
84. Raaymakers TW. Aneurysms in relatives of patients with subarachnoid hemorrhage: Frequency and risk factors. MARS Study Group. Magnetic resonance angiography in relatives of patients with Subarachnoid hemorrhage. *Neurology* 1999; 53:982-88.
85. Kasuya H, Onda H, Takeshita M, Hori T, Takakura K. Clinical features of intracranial aneurysms in siblings. *Neurosurgery* 2000; 46:1301-05.
86. Foroud T, Sauerbeck L, Brown R, Anderson C, Woo D, Kleindorfer D, et al. Genome screen to detect linkage to intracranial aneurysm susceptibility genes: the Familial Intracranial Aneurysm (FIA) study. *Stroke* 2008; 39:1434-40.
87. Bromberg JE, Rinkel GJ, Algra A, Greebe P, van Duyn CM, Hasan D, et al. Subarachnoid haemorrhage in first and second degree relatives of patients with subarachnoid haemorrhage. *BMJ* 1995a;311:288-89.
88. Bromberg JE, Rinkel GJ, Algra A, van Duyn CM, Greebe P, Ramos LM, et al. Familial subarachnoid hemorrhage: distinctive features and patterns of inheritance. *Ann Neurol* 1995b; 38: 929-34.
89. Teasdale GM, Wardlaw JM, White PM, Murray G, Teasdale EM, Easton V, et al. The familial risk of subarachnoid haemorrhage. *Brain* 2005; 128: 1677-85.
90. Nakagawa T, Hashi K, Kurokawa Y, Yamamura A. Family history of subarachnoid hemorrhage and the incidence of asymptomatic, unruptured cerebral aneurysms. *J Neurosurg* 1999;91:391-95.

91. Kissela BM, Sauerbeck L, Woo D, Khoury J, Carrozzella J, Pancioli A, et al. Subarachnoid hemorrhage: A preventable disease with a heritable component. *Stroke* 2002; 33:1321-26.
92. Brown RD Jr, Huston J, Hornung R, Foroud T, Kallmes DF, et al. Screening for brain aneurysm in the Familial Intracranial Aneurysm study: frequency and predictors of lesion detection. *J Neurosurg* 2008; 108:1132-38.
93. Schievink WI, Wijdicks EF, Parisi JE, Piepgras DG, Whisnant JP. Sudden death from aneurysmal subarachnoid hemorrhage. *Neurology* 1995b;45: 871-74.
94. Wang PS, Longstreth Jr WT, Koepsell TD. Subarachnoid hemorrhage and family history. A population-based case-control study. *Arch Neurol* 1995a;52:202-4.
95. Rinkel GJ. Intracranial aneurysm screening: indications and advice for practice. *Lancet Neurol* 2005; 4:122-28.
96. Rinkel GJ, Djibuti M, Algra A, van Gijn J. Prevalence and risk of rupture of intracranial aneurysms: a systematic review. *Stroke* 1998;29:251-256.
97. De Brackeleer M, Perusse L, Cantin L, Bouchard JM, Mathieu J. A study of inbreeding and kinship in intracranial aneurysms in the Saguenay Lac-Saint-Jean region (Quebec, Canada). *Ann Hum Genet* 1996;60:99-104.
98. Kirkpatrick PJ, McConnell RS. Screening for familial intracranial aneurysms. *BMJ* 1999;319:1512-13.
99. Schievink WI. Genetics of intracranial aneurysms. *Neurosurgery* 1997a;40:651-662.
100. Schievink WI. Intracranial aneurysms. *N Engl J Med* 1997b;336:28-40.
101. Mackey J, Brown RD Jr, Moomaw CJ, Sauerbeck L, Hornung R, Gandhi D, et al. Unruptured intracranial aneurysms in the Familial Intracranial Aneurysm and

- International Study of Unruptured Intracranial Aneurysms cohorts: differences in multiplicity and location. *J Neurosurg* 2012;117:60-4.
102. Lozano AM, Leblanc R. Familial intracranial aneurysms. *J Neurosurg* 1987; 66:522-28.
103. Onda H, Kasuya H, Yoneyama T, Takakura K, Hori T, Takeda J, et al. Genomewide-linkage and haplotype- association studies map intracranial aneurysm to chromosome 7q11. *Am J Hum Genet* 2001; 69: 804-819.
104. Pannu H, Kim DH, Guo D, King TM, Van Ginhoven G, Chin T, Chang K, Qi Y, Shete S, Milewicz DM. The role of MMP-2 and MMP-9 polymorphisms in sporadic intracranial aneurysms. *J Neurosurg* 2006;105:418-23.
105. Kuivaniemi H, Prockop DJ, Wu Y, Madhatheri SL, Kleinert C, Earley JJ, et al. Exclusion of mutations in the gene for type III collagen (COL3A1) as a common cause of intracranial aneurysms or cervical artery dissections: results from sequence analysis of the coding sequences of type III collagen from 55 unrelated patients. *Neurology* 1993; 43:2652-58.
106. Dai B, Liu T, Zhang B, Zhang X, Wang Z. The polymorphism for endothelial nitric oxide synthase gene, the level of nitric oxide and the risk for pre-eclampsia: a meta-analysis. *Gene* 2013;519:187-93.
107. Ignarro LJ, Cirino G, Casini A, Napoli C. Nitric oxide as a signaling molecule in the vascular system: an overview. *J Cardiovasc Pharmacol* 1999;34:879-86.
108. Chen W, Srinivasan SR, Elkasabany A, Ellsworth DL, Boerwinkle E, Berenson GS. Combined effects of endothelial nitric oxide synthase gene polymorphism (G894T) and insulin resistance status on blood pressure and familial risk of

- hypertension in young adults: the Bogalusa Heart Study. *Am J Hypertens* 2001;14:1046–52.
109. Chen W, Srinivasan SR, Bond MG, Tang R, Urbina EM, Li S, et al. Nitric oxide synthase gene polymorphism (G894T) influences arterial stiffness in adults: the Bogalusa Heart Study. *Am J Hypertens* 2004a;17: 553–59.
110. Chen W, Srinivasan SR, Li S, Boerwinkle E, Berenson GS. Gender specific influence of NO synthase gene on blood pressure since childhood: the Bogalusa Heart Study. *Hypertension* 2004b;44:668 –73.
111. Khurana VG, Meissner I, Meyer FB. Update on genetic evidence for rupture-prone compared with rupture-resistant intracranial saccular aneurysms. *Neurosurg Focus* 2004a;17:E7.
112. Napoli C, de Nigris F, Williams-Ignarro S, Pignalosa O, Sica V, Ignarro LJ. Nitric oxide and atherosclerosis: an update. *Nitric Oxide* 2006;15:265-79.
113. Sadamitsu D, Kuroda Y, Nagamitsu T, Tsuruta R, Inoue T, Ueda T, et al. Cerebrospinal fluid and plasma concentrations of nitric oxide metabolites in postoperative patients with subarachnoid hemorrhage. *Crit Care Med* 2001; 29: 77-9.
114. Bhuiyan AR, Chen W, Srinivasan SR, Rice J, Mock N, Tang R, et al. Influence of nitric oxide synthase gene polymorphism (G894T) on carotid artery intima-media thickness in adults: the Bogalusa Heart Study. *J Am Soc Hypertens* 2007;1:362-68.
115. Mayer O Jr, Filipovský J, Pesta M, Cífková R, Dolejšová M, Simon J. The interaction of endothelial nitric oxide synthase polymorphism and current smoking in terms of increased arterial stiffness. *Physiol Res* 2010;59:529-36.

116. Paik D, Tilson MD. Neovascularization in the abdominal aortic aneurysm. Endothelial nitric oxide synthase, nitric oxide, and elastolysis. *Ann N Y Acad Sci* 1996;800:277.
117. Dosenko VE, Zagoriy VY, Haytovich NV, Gordok OA, Moibenko AA. Allelic polymorphism of endothelial NO-synthase gene and its functional manifestations. *Acta Biochim Pol* 2006; 53:299-302.
118. Johanning JM, Armstrong PJ, Franklin DP, Han DC, Carey DJ, Elmore JR. Nitric oxide in experimental aneurysm formation: early events and consequences of nitric oxide inhibition. *Ann Vasc Surg* 2002;16:65-72.
119. Mineharu Y, Inoue K, Inoue S, Yamada S, Nozaki K, Takenaka K, et al. Association analysis of common variants of ELN, NOS2A, APOE and ACE2 to intracranial aneurysm. *Stroke* 2006; 37:1189-94.
120. Kim TG, Kim NK, Baek MJ, Huh R, Chung SS, Choi JU, et al. The relationships between endothelial nitric oxide synthase polymorphisms and the formation of intracranial aneurysms in the Korean population. *Neurosurg Focus* 2011;30:E23.
121. Marsden PA, Heng HH, Scherer SW, Stewart RJ, Hall AV, Shi XM, et al. Structure and chromosomal localization of the human constitutive endothelial nitric oxide synthase gene. *J Biol Chem* 1993;268:17478–488.
122. Wattanapitayakul SK, Mihm MJ, Young AP, Bauer JA. Therapeutic implications of human endothelial nitric oxide synthase gene polymorphism. *Trends Pharmacol Sci* 2001; 22: 361-8.
123. Veldman BA, SpieringW, Doevendans PA, Vervoort G, Kroon AA, de Leeuw PW, Smits P. The Glu298Asp polymorphism of the NOS 3 gene as a determinant of the baseline production of nitric oxide. *J Hypertens* 2002;20: 2023–27.

124. Saracini C, Bolli P, Sticchi E, Pratesi G, Pulli R, Sofi F, Pratesi C, Gensini GF, Abbate R, Giusti B. Polymorphisms of genes involved in extracellular matrix remodeling and abdominal aortic aneurysm. *J Vasc Surg* 2012;55:171-79.
125. Li DY, Brooke B, Davis EC, Mecham RP, Sorensen LK, Boak BB, et al. Elastin is an essential determinant of arterial morphogenesis. *Nature* 1998; 393: 276-80.
126. Sauvage M, Hinglais N, Mandet C, Badier C, Deslandes F, Michel JB, Jacob MP. Localization of elastin mRNA and TGF- $\beta$ 1 in rat aorta and caudal artery as a function of age. *Cell Tissue Res* 1998;291:305-14.
127. Fleenor BS, Seals DR, Zigler ML, Sindler AL. Superoxide-lowering therapy with TEMPOL reverses arterial dysfunction with aging in mice. *Aging Cell* 2012b;11:269-76.
128. Fleenor BS, Sindler AL, Eng JS, Nair DP, Dodson RB, Seals DR. Sodium nitrite de-stiffening of large elastic arteries with aging: Role of normalization of advanced glycation end-products. *Exp Gerontol* 2012c;47:588-94.
129. Ewart AK, Morris CA, Atkinson D, Jin W, Sternes K, Spallone P, et al. Hemizyosity at the elastin locus in a developmental disorder, Williams syndrome. *Nat Genet.* 1993a;5:11-6.
130. Ewart AK, Morris CA, Ensing GJ, Loker J, Moore C, Leppert M, et al. A human vascular disorder, supraaortic stenosis, maps to chromosome 7. *Proc Natl Acad Sci U S A* 1993b; 90:3226-30.
131. Raybould MC, Birley AJ, Hulten M. A Bgl I polymorphism in the human elastin gene (ELN). *Clin Genet* 1994a;46:212-13.
132. Raybould MC, Birley AJ, Hulten M. Two new polymorphisms in the human elastin gene (ELN). *Hum Genet* 1994b;93:475-76.

133. Ruigrok YM, Rinkel GJE. Genetics of intracranial aneurysms. *Stroke* 2008;39:1049–55.
134. Farnham JM, Camp NJ, Neuhausen SL, Tsuruda J, Parker D, MacDonald J, et al. Confirmation of chromosome 7q11 locus for predisposition to intracranial aneurysm. *Hum Genet* 2004; 114:250-55.
135. Olson JM, Vongpunsawad S, Kuivaniemi H, Ronkainen A, Hernesniemi J, Ryyänen M, Kim LL, Tromp G. Search for intracranial aneurysm susceptibility gene(s) using Finnish families. *BMC Med Genet* 2002; 3: 7.
136. Tromp G, Christiano A, Goldstein N, Indik, Z, Boyd C, Rosenbloom J. A to G polymorphism in ELN gene. *Nucleic Acids Res* 1991;19:4314-15.
137. Hanon O, Luong V, Mourad JJ, Bortolotto LA, Jeunemaitre X, Girerd X. Aging, Carotid Artery Distensibility, and the Ser422Gly Elastin Gene Polymorphism in Humans. *Hypertension* 2001; 38:1185-89.
138. Ogata T, Shibamura H, Tromp G, Sinha M, Goddard KA, Sakalihasan N, et al. Genetic analysis of polymorphisms in biologically relevant candidate genes in patients with abdominal aortic aneurysms. *J Vasc Surg* 2005;41:1036-42.
139. Yamada S, Utsunomiya M, Inoue K, Nozaki K, Miyamoto S, Hashimoto N, et al. Absence of linkage of familial intracranial aneurysms to 7q11 in highly aggregated Japanese families. *Stroke* 2003;34: 892-900.
140. Hofer A, Hermans M, Kubassek N, Sitzer M, Funke H, Stogbauer F, et al. Elastin Polymorphism Haplotype and Intracranial Aneurysms are not associated in Central Europe. *Stroke* 2003;34: 1207-11.
141. Hanahan D. Signaling vascular morphogenesis and maintenance. *Science* 1997; 277:48-50.



142. Gougos A, Letarte M. Primary structure of endoglin, an RGD-containing glycoprotein of human endothelial cells. *J Biol Chem* 1990;265:8361–64.
143. Wang XF, Lin HY, Ng-Eaton E, Downward J, Lodish HF, Weinberg RA. Expression cloning and characterization of the TGF- $\beta$  type III receptor. *Cell* 1991;67:797–805.
144. Wong SH, Hamel L, Chevalier S, Philip A. Endoglin expression on human microvascular endothelial cells association with betaglycan and formation of higher order complexes with TGF- $\beta$  signalling receptors. *Eur J Biochem* 2000;267:5550–60.
145. Joo SP, Kim TS, Lee IK, Lee JK, Seo BR, Kim JH, et al. The role of collagen type I alpha2 polymorphisms: intracranial aneurysms in Koreans. *Surg Neurol* 2009; 72: 48-53.
146. López-Novoa JM, Bernabeu C. The physiological role of endoglin in the cardiovascular system. *Am J Physiol Heart Circ Physiol* 2010; 299: H959-74.
147. Morris E, Chrobak I, Bujor A, Hant F, Mummery C, Ten Dijke P, Trojanowska M. Endoglin promotes TGF- $\beta$ /Smad1 signaling in scleroderma fibroblasts. *J Cell Physiol* 2011; 226: 3340-48.
148. Calabrò L, Fonsatti E, Bellomo G, Alonci A, Colizzi F, Sigalotti L, et al. Differential levels of soluble endoglin (CD105) in myeloid malignancies. *J Cell Physiol* 2003;194:171–75.
149. Levine RJ, Lam C, Qian C, Yu KF, Maynard SE, Sachs BP, et al. Soluble endoglin and other circulating antiangiogenic factors in preeclampsia. *N Engl J Med* 2006;355:992–1005.

- 
150. Venkatesha S, Toporsian M, Lam C, Hanai J, Mammoto T, Kim YM, et al. Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 2006;2:642–49.
  151. Fonsatti E, Jekunen AP, Kairemo KJ, Coral S, Snellman M, Nicotra MR, et al. Endoglin is a suitable target for efficient imaging of solid tumors: in vivo evidence in a canine mammary carcinoma model. *Clin Cancer* 2000;Res 6:2037–43.
  152. Sánchez-Elsner T, Botella LM, Velasco B, Langa C, Bernabéu C. Endoglin expression is regulated by transcriptional cooperation between the hypoxia and transforming growth factor- $\beta$  pathways. *J Biol Chem* 2002;277:43799–808.
  153. van Laake LW, van den Driesche S, Post S, Feijen A, Jansen MA, Driessens MH, et al. Endoglin has a crucial role in blood cell-mediated vascular repair. *Circulation* 2006;114:2288–97.
  154. Torsney E, Charlton R, Parums D, Collis M, Arthur HM. Inducible expression of human endoglin during inflammation and wound healing in vivo. *Inflamm Res* 2000;51:464–70.
  155. Jonker L, Arthur HM. Endoglin expression in early development is associated with vasculogenesis and angiogenesis. *Mech Dev* 2002;110:193–96.
  156. ten Dijke P, Goumans MJ, Pardali E. Endoglin in angiogenesis and vascular diseases. *Angiogenesis* 2008;11:79–89.
  157. Arthur HM, Ure J, Smith AJ, Renforth G, Wilson DI, Torsney E, et al. Endoglin, an ancillary TGF $\beta$  receptor, is required for extraembryonic angiogenesis and plays a key role in heart development. *Dev Biol* 2000;217:42–53.

158. Ma X, Labinaz M, Goldstein J, Miller H, Keon WJ, Letarte M, et al. Endoglin is overexpressed after arterial injury and is required for transforming growth factor-beta-induced inhibition of smooth muscle cell migration. *Arterioscler Thromb Vasc Biol* 2000; 20: 2546-52.
159. Mancini ML, Terzic A, Conley B, Oxburgh LH, Nicola T, Vary C. Endoglin Plays Distinct Roles in Vascular Smooth Muscle Cell Recruitment and Regulation of Arteriovenous Identity During Angiogenesis. *Dev Dyn* 2009;238:2479-93.
160. Wong G, Knuckey NW and Gubbay SS. Subarachnoid haemorrhage in children caused by cerebral tumour. *J Neurol Neurosurg Psychiatry* 1983; 46: 449–50.
161. Botella LM, Sanz-Rodriguez F, Sanchez-Elsner T, Langa C, Ramirez JR, Vary C, et al. Lumican is down-regulated in cells expressing endoglin. Evidence for an inverse corelationship between endoglin and lumican expression. *Matrix Biol* 2004; 22: 561-72.
162. Onda H, Kasuya H, Yoneyama T, Hori T, Nakajima T, Inoue I. Endoglin is not a major susceptibility gene for intracranial aneurysm among Japanese. *Stroke* 2003; 34: 1640-44.
163. McColgan P, Thant KZ, Sharma P. The genetics of sporadic ruptured and unruptured intracranial aneurysms: a genetic meta-analysis of 8 genes and 13 polymorphisms in approximately 20,000 individuals. *J Neurosurg* 2010; 112:714–21.
164. Stehbens WE. *Pathology of the cerebral blood vessels*. St Louis: Mosby, 1972.
165. Gläsker S, Schatlo B, Klingler JH, Braun V, Spangenberg P, Kim IS, et al. Associations of Collagen Type I  $\alpha 2$  Polymorphisms with the Presence of

- Intracranial Aneurysms in Patients from Germany. *J Stroke Cerebrovasc Dis.* 2013 Jun 22.
166. Dagleish R. The human type I collagen mutation database. *Nucleic Acids Res* 1997; 25:181-87.
167. Engel J, Prockop DJ. The zipper-like folding of collagen triple helices and the effects of mutations that disrupt the zipper. *Annu Rev Biophys Biophys Chem* 1991;20:137-52.
168. Prockop DJ, Kivirikko KI. Collagens: molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem* 1995; 64: 403-34.
169. Ikonomidis JS, Jones JA, Barbour JR, Stroud RE, Clark LL, Kaplan BS, et al. Expression of matrix metalloproteinases and endogenous inhibitors within ascending aortic aneurysms of patients with Marfan syndrome. *Circulation* 2006;114:I365-70.
170. Zuo C, Wen F, Li M, Zhang X, Chen H, Wu K, Zeng R. COL1A2 polymorphic markers confer an increased risk of neovascular age-related macular degeneration in a Han Chinese population. *Mol Vis.* 2012;18:1787-93.
171. Niederreither K, D'Souza RN, de Crombrughe B. Minimal DNA sequences that control the cell lineage-specific expression of the pro alpha 2(I) collagen promoter in transgenic mice. *J Cell Biol* 1992; 119:1361-70.
172. Ponticos M, Abraham D, Alexakis C, Lu QL, Black C, Partridge T, Bou-Gharios G. Colla2 enhancer regulates collagen activity during development and in adult tissue repair. *Matrix Biol* 2004; 22:619-28.

- 
173. Lindahl K, Rubin CJ, Brandstrom H, Karlsson MK, Holmberg A, Ohlsson C, et al. Heterozygosity for a coding SNP in COL1A2 confers a lower BMD and an increased stroke risk. *Biochem Biophys Res Commun* 2009; 384:501-5.
174. Allt G, Lawrenson JG. Pericytes cell biology and pathology. *Cells Tissues Organs* 2001;169:1-11.
175. von Tell D, Armulik A, Betsholtz C. Pericytes and vascular stability. *Exp Cell Res* 2006;312:623-29.
176. Yoneyama T, Kasuya H, Onda H, Akagawa H, Hashiguchi K, Nakajima T, et al. Collagen type I alpha2 (*COL1A2*) is the susceptible gene for intracranial aneurysms. *Stroke* 2004;5:443-48.
177. Roos YB, Pals G, Struycken PM, Rinkel GJ, Limburg M, Pronk JC, van den Berg JS, Luijten JA, Pearson PL, Vermeulen M, Westerveld A. Genome-wide linkage in a large Dutch consanguineous family maps a locus for intracranial aneurysms to chromosome 2p13. *Stroke* 2004; 35:2276-81.
178. Roder C, Kasuya H, Harati A, Tatagiba M, Inoue I, Krischek B. Meta-analysis of microarray gene expression studies on intracranial aneurysms. *Neuroscience* 2012;201:105-13.
179. Zhu Y, Li W, Ge M, Xu S, Zhao G, Wang H, et al. Polymorphism rs42524 of *COL1A2* and sporadic intracranial aneurysms in the Chinese population. *J Neurosurg* 2008;109:1060-64.
180. Muhlhauser J, Merrill MJ, Pili R, Maeda H, Bacic M, Bewig B, et al. VEGF165 expressed by a replication-deficient recombinant adenovirus vector induces angiogenesis in vivo. *Circ Res* 1995; 77: 1077-86.

181. Safi J, Gloe TR, Riccioni T, Kovesdi I, Capogrossi MC. Gene therapy with angiogenic factors: a new potential approach to the treatment of ischemic diseases. *J Mol Cell Cardiol* 1997; 29: 2311–25.
182. Jiang Y, Liang G, Wang L, Jiang J, Du G, Huang Y. Association between vascular endothelial growth factor +936 C/T gene polymorphism and age-related macular degeneration *J Int Med Res* 2013;41:317-24.
183. Renner W, Kotschan S, Hoffmann C, Obermayer-Pietsch B, Pilger E. A common 936 C/T mutation in the gene for vascular endothelial growth factor is associated with vascular endothelial growth factor plasma levels. *J Vasc Res* 2000; 37: 443-8.
184. Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J. Vascular-specific growth factors and blood vessel formation. *Nature* 2000;407(6801):242-8.
185. Bikfalvi A. Recent developments in the inhibition of angiogenesis: examples from studies on platelet factor-4 and the VEGF/VEGFR system. *Biochem Pharmacol* 2004;68(6):1017-21.
187. Kilic T, Sohrabifar M, Kurtkaya O, Yildirim O, Elmaci I, Gunel M, et al. Expression of structural proteins and angiogenic factors in normal arterial and unruptured and ruptured aneurysm walls. *Neurosurg* 2005;57:997–1007.
188. Ferrara N, Gerber HP, Le Couter J. The biology of VEGF and its receptors. *Nat Med* 2003; 9: 669–76.
189. Ferrara N, Gerber HP. The role of vascular endothelial growth factor in angiogenesis. *Acta Haematol* 2001; 106:148-56.
190. Gonçalves FM, Martins-Oliveira A, Speciali JG, Izidoro-Toledo TC, Luizon MR, Dach F, et al. Vascular endothelial growth factor genetic polymorphisms and haplotypes in women with migraine. *DNA Cell Biol* 2010; 29: 357–62.

191. Houck KA, Ferrara N, Winer J, Cachianes G, Li B, Leung DW. The vascular endothelial growth factor family: identification of a fourth molecular species and characterization of alternative splicing of RNA. *Mol Endocrinol* 1991; 5:1806-14.
192. Watson CJ, Webb NJ, Bottomley MJ, Brenchley PE. Identification of polymorphisms within the vascular endothelial growth factor (VEGF) gene: correlation with variation in VEGF protein production. *Cytokine* 2000;12:1232-5.
193. Awata T, Inoue K, Kurihara S, Ohkubo T, Watanabe M, Inukai K, et al. A common polymorphism in the 5'-untranslated region of the VEGF gene is associated with diabetic retinopathy in type 2 diabetes. *Diabetes* 2002;51:1635-9.
194. Papazoglou D, Galazios G, Papatheodorou K, Liberis V, Papanas N, Maltezos E, ET AL. Vascular endothelial growth factor gene polymorphisms and idiopathic recurrent pregnancy loss. *Fertil Steril* 2005;83:959-63.
195. Samli H, Demir BÇ, Ozgöz A, Atalay MA, Uncu G. Vascular endothelial growth factor gene 1154 G/A, 2578 C/A, 460 C/T, 936 C/T polymorphisms and association with recurrent pregnancy losses. *Genet Mol Res* 2012;11:4739-45.
196. He Y, Ni J, Chen S, Jiang Y, Jia S, Gao Y. The vascular endothelial growth factor-2549 insertion/deletion polymorphism is not associated with susceptibility to hepatocellular carcinoma in Chinese. *DNA Cell Biol* 2010;29:393-96.
197. Belo VA, Souza-Costa DC, Luizon MR, Izidoro-Toledo TC, Lanna CM, Pinheiro LC, et al. Vascular endothelial growth factor haplotypes associated with childhood obesity. *DNA Cell Biol* 2011; 30: 709-714.
198. Mittal RD, Srivastava P, Singh V, Jaiswal P, Kapoor R. Association of common variants of vascular endothelial growth factor and interleukin-18 genes with

- allograft survival in renal transplant recipients of North India. *DNA Cell Biol* 2011; 30: 309–15.
199. Lu Y, Shi Y, Xue C, Yin J, Huang Z. Pooled-analysis of the associations between three polymorphisms in the VEGF gene and age-related macular degeneration. *Mol Biol Rep* 2012; 39: 6547–53.
200. Qu Y, Dai H, Zhou F, Zhang X, Xu X, Zhang X, et al. Vascular endothelial growth factor gene polymorphisms and risk of neovascular age-related macular degeneration in a Chinese cohort. *Ophthalmic Res* 2011;45:142–48.
201. Ajaz S, Khaliq S, Abid A, Hassan AS, Hashmi A, Sultan G, et al. Association of a single-nucleotide polymorphism in the promoter region of the VEGF gene with the risk of renal cell carcinoma. *Genet Test Mol Biomarkers* 2011;15:653–57.
202. Zhang L, Zhang G, Wang P, Gong J, Cao Y, Tang L. Association of vascular endothelial growth factor -2578C/A gene polymorphism in Chinese patients with colon cancer. *Genet Test Mol Biomarkers* 2011; 15: 117–21.
203. Skirgaudas M, Awad IA, Kim J, Rothbart D, Criscuolo G. Expression of angiogenesis factors and selected vascular wall matrix proteins in intracranial saccular aneurysms. *Neurosurgery* 1996;39:537-45.
204. Maderna E, Corsini E, Franzini A, Giombini S, Pollo B, Broggi G, et al. Expression of vascular endothelial growth factor receptor-1/-2 and nitric oxide in unruptured intracranial aneurysms. *Neurol Sci* 2010; 31:617–23.
186. Brooks PC. Role of integrin in angiogenesis. *Eur J Cancer* 1996;32A:2423–29.
205. Zhou LF. Cerebral aneurysm. Zhou LF ed. *Modern Neurosurgery*. Shanghai: Fudan University Publishing House; 2001:816-20.



206. Fukuda S, Hashimoto N, Naritomi H, Nagata I, Nozaki K, Kondo S, et al. Prevention of rat cerebral aneurysm formation by inhibition of nitric oxide synthase. *Circulation* 2000;101: 2532–38.
207. Tateshima S, Murayama Y, Villablanca JP, Morino T, Takahashi H, Yamauchi T, et al. Intraaneurysmal flow dynamics study featuring an acrylic aneurysm model manufactured using a computerized tomography angiogram as a mold. *J Neurosurg* 2001;95:1020-27.
208. Morimoto M, Miyamoto S, Mizoguchi A, Kume N, Kita T, Hashimoto N. Mouse model of cerebral aneurysm experimental induction by renal hypertension and local hemodynamic changes. *Stroke* 2002;33:1911-15.
209. Juvela S, Hillbom M, Numminen H, Koskinen P. Cigarette smoking and alcohol consumption as risk factors for aneurysmal subarachnoid hemorrhage. *Stroke* 1993a; 24: 639-46.
210. Juvela S, Porras M, Heiskanen O. Natural history of unruptured intracranial aneurysms: a long-term follow-up study. *J Neurosurg* 1993b;79:174-82.
211. Ingall T, Asplund K, Mähönen M, Bonita R. A multinational comparison of subarachnoid hemorrhage epidemiology in the WHO MONICA stroke study. *Stroke* 2000; 31: 1054-61.
212. Ellamushi HE, Grieve JP, Jager HR, Kitchen ND. Risk factors for the formation of multiple intracranial aneurysms. *J Neurosurg* 2001; 94:728-32.
213. Broderick JP, Viscoli CM, Brott T, Kernan WN, Brass LM, Feldmann E, et al. Major risk factors for aneurysmal subarachnoid hemorrhage in the young are modifiable. *Stroke* 2003;34:1375-81.

214. Juvela S. Prehemorrhage risk factors for fatal intracranial aneurysm rupture. *Stroke* 2003; 34:1852-57
216. Ohkuma H, Tabata H, Suzuki S, Islam MS. Risk factors for aneurysmal subarachnoid hemorrhage in Aomori, Japan. *Stroke* 2003;34:96-100.
217. Anderson CS, Feigin V, Bennett D, Lin RB, Hankey G, Jamrozik K, Australasian Cooperative Research on Subarachnoid Hemorrhage Study (ACROSS) Group. Active and passive smoking and the risk of subarachnoid hemorrhage: an international population-based case- control study. *Stroke* 2004; 35: 633-37.
218. Feigin VL, Parag V, Lawes CM, Rodgers A, Suh I, Woodward M, et al. Smoking and elevated blood pressure are the most important risk factors for subarachnoid hemorrhage in the Asia-Pacific region: an overview of 26 cohorts involving 306,620 participants. *Stroke* 2005a; 36:1360-65.
215. Wiebers DO, Whisnant JP, Huston J 3rd, Meissner I, Brown RD Jr, Piepgras DG, et al. Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment. *Lancet* 2003; 362:103-10.
219. Feigin VL, Rinkel GJ, Lawes CM, Algra A, Bennett DA, van Gijn J, et al. Risk factors for subarachnoid hemorrhage: an updated systematic review of epidemiological studies. *Stroke* 2005b;36: 2773-80.
220. Kawachi I, Colditz GA, Stampfer MJ, Willett WC, Manson JE, Rosner B, et al. Smoking cessation and decreased risk of stroke in women. *JAMA* 1993;269:232–36.
221. Qureshi AI, Suarez JJ, Parekh PD, Sung G, Geocadin R, Bhardwaj A, Tamargo RJ, Ulatowski JA. Risk factors for multiple intracranial aneurysms. *Neurosurgery* 1998;43:22-6.

- 
222. Juvela S, Porras M, Poussa K. Natural history of unruptured intracranial aneurysms: probability of and risk factors for aneurysm rupture. *J Neurosurg* 2000;93:379-87.
223. Taylor CL, Yuan Z, Selman WR, Ratcheson RA, Rimm AA. Cerebral arterial aneurysm formation and rupture in 20,767 elderly patients: hypertension and other risk factors. *J Neurosurg* 1995;83:812-9.
224. Qureshi AI, Suri MF, Yahia AM, Suarez JJ, Guterman LR, Hopkins LN, et al. Risk factors for subarachnoid hemorrhage. *Neurosurg* 2001;49:607-13.
225. Owens, WB. Blood Pressure Control in Acute Cerebrovascular Disease. *J Clin Hypertens (Greenwich)* 2011;13:205-11
226. Juvela S, Poussa K, Porras M. Factors affecting formation and growth of intracranial aneurysms: a long-term follow-up study. *Stroke* 2001; 32:485-91.
227. Khurana VG, Meissner I, Sohni YR, Bamlet WR, McClelland RL, Cunningham JM, et al. The presence of tandem endothelial nitric oxide synthase gene polymorphisms identifying brain aneurysms more prone to rupture. *J Neurosurg* 2005;102:526–31.
228. Woo D, Khoury J, Haverbusch MM, Sekar P, Flaherty ML, Kleindorfer DO, et al. Smoking and family history and risk of aneurysmal subarachnoid hemorrhage. *Neurology* 2009;72:69-72.
229. Deka R, Koller DL, Lai D, Indugula SR, Sun G, Woo D, et al. The relationship between smoking and replicated sequence variants on chromosomes 8 and 9 with familial intracranial aneurysm. *Stroke* 2010;41:1132-37.
230. Chrysohoou C, Panagiotakos DB, Pitsavos C, Antoniadis C, Skoumas J, Brown M, Stefanadis C. Evidence for association between endothelial nitric oxide synthase

- gene polymorphism (G894T) and inflammatory markers: the ATTICA study. *Am Heart J* 2004;148:733-38.
231. Siasos G, Tousoulis D, Vlachopoulos C, Antoniades C, Stefanadi E, Ioakeimidis N, et al. Short-term treatment with L-arginine prevents the smoking-induced impairment of endothelial function and vascular elastic properties in young individuals. *Int J Cardiol* 2008;126:394-9.
232. Chalouhi N, Ali MS, Jabbour PM, Tjoumakaris SI, Gonzalez LF, Rosenwasser RH, et al. Biology of intracranial aneurysms: role of inflammation. *J Cereb Blood Flow Metab* 2012;32:1659-76.
233. Mayhan WG, Patel KP. Effect of nicotine on endothelium-dependent arteriolar dilatation in vivo. *Am J Physiol* 1997;272:H2337-42.
234. Mayhan WG, Sharpe GM, Anding P. Agonist-induced release of nitric oxide during acute exposure to nicotine. *Life Sci* 1999;65:1829-37.
235. Koide M, Nishizawa S, Yamamoto S, Yamaguchi M, Namba H, Terakawa S. Nicotine exposure, mimicked smoking, directly and indirectly enhanced protein kinase C activity in isolated canine basilar artery, resulting in enhancement of arterial contraction. *J Cereb Blood Flow Metab* 2005;25:292-301.
236. Gerzanich V, Zhang F, West GA, Simard JM. Chronic nicotine alters NO signaling of Ca(2+) channels in cerebral arterioles. *Circ Res* 2001;88:359-65.
237. Dumont AS, Kassell NF. Editorial comment - parity and risk of subarachnoid hemorrhage: an emerging association. *Stroke* 2004;35:32-3.
238. Fujita J, Skold CM, Daughton DM, Ertl RF, Takahara J, Rennard SI. Modulation of elastase binding to elastin by human alveolar macrophage-derived lipids. *Am J Respir Crit Care Med* 1999; 160: 802-07.

239. Shinton R, Beevers G. Meta-analysis of relation between cigarette smoking and stroke. *Br Med J* 1989;298:789-94.
240. Weir BK, Kongable GL, Kassell NF, Schultz JR, Truskowski LL, Sigrest A. Cigarette smoking as a cause of aneurysmal subarachnoid hemorrhage and risk for vasospasm: a report of the Cooperative Aneurysm Study. *J Neurosurg* 1998;89:405-11.
241. Qureshi AI, Sung GY, Suri MFK, Straw RN, Guterman LR, Hopkins LN. Factors associated with aneurysm size in patients with subarachnoid hemorrhage: effect of smoking and aneurysm location. *Neurosurgery* 2000;46:44-50.
242. World Health Organization, 2013 - <http://www.who.int/en/>
243. Schmidt MI, Duncan BB, Hoffmann JF, Moura Ld, Malta DC, Carvalho RM. Prevalence of diabetes and hypertension based on self-reported morbidity survey, Brazil, 2006. *Rev Saude Publica*. 2009 Nov;43 Suppl 2:74-82.
244. De la Monte SM, Moore GW, Monk MA, Hutchins GM. Risk factors for the development and rupture of intracranial berry aneurysms. *Am J Med* 1985;78:957-64.
245. Lindström J, Tuomilehto J. The diabetes risk score: a practical tool to predict type 2 diabetes risk. *Diabetes Care* 2003;26:725–31.
246. Lyssenko V, Jonsson A, Almgren P, Pulizzi N, Isomaa B, Tuomi T, et al. Clinical risk factors, DNA variants, and the development of type 2 diabetes. *N Engl J Med* 2008;359:2220–32.
247. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2011;34(Suppl. 1):S62–S69.

- 
249. Almdal T, Scharling H, Jensen JS, Vestergaard H. The independent effect of type 2 diabetes mellitus on ischemic heart disease, stroke, and death: a population-based study of 13,000 men and women with 20 years of follow-up. *Arch Intern Med* 2004;164:1422–26.
248. Nolan CJ, Damm P, Prentki M. Type 2 diabetes across generations: from pathophysiology to prevention and management. *Lancet* 2011;378:169–81.
250. O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010;376:112–23.
251. Emerging Risk Factors Collaboration, Sarwar N, Gao P, Seshasai SR, Gobin R, Kaptoge S, et al. Emerging Risk Factors Collaboration. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet* 2010;375:2215–22.
252. Okamoto K, Horisawa R, Ohno Y. The relationships of gender, cigarette smoking, and hypertension with the risk of aneurysmal subarachnoid hemorrhage: a case-control study in Nagoya, Japan. *Ann Epidemiol* 2005;15:744–48.
253. Koshy L, Easwer HV, Premkumar S, Alapatt JP, Pillai AM, Nair S, et al. Risk factors for aneurysmal subarachnoid hemorrhage in an Indian population. *Cerebrovasc Dis* 2010;29:268–74.
254. Cui R, Iso H, Yamagishi K, Saito I, Kokubo Y, Inoue M, et al. Diabetes mellitus and risk of stroke and its subtypes among Japanese: the Japan public health center study. *Stroke* 2011;42:2611–14.

255. Shiue I, Arima H, Hankey GJ, Anderson CS, ACROSS Group. Modifiable lifestyle behaviours account for most cases of subarachnoid haemorrhage: a populationbased case-control study in Australasia. *J Neurol Sci* 2012;313:92-4.
256. Lindgren AE, Kurki MI, Riihinen A, Koivisto T, Ronkainen A, Rinne J, et al. Type 2 diabetes and risk of rupture of saccular intracranial aneurysm in eastern Finland. *Diabetes Care* 2013;36:2020-26.
257. Keenan NL, Rosendorf KA, Centers for Disease Control and Prevention (CDC). Prevalence of hypertension and controlled hypertension - United States, 2005-2008. *MMWR Surveill Summ* 2011;60 Suppl:94-7.
258. Ferreira SR, Moura EC, Malta DC, Sarno F. Frequency of arterial hypertension and associated factors: Brazil, 2006. *Rev Saude Publica* 2009;43 Suppl 2:98-106.
259. Nitrini R, Caramelli P, Herrera E Jr, de Castro I, Bahia VS, Anghinah R, et al. Mortality from dementia in a community-dwelling Brazilian population. *Int J Geriatr Psychiatry* 2005;20:247-53.
260. Connolly ES Jr, Choudhri TF, Mack WJ, Mocco J, Spinks TJ, Slosberg J, et al. Influence of smoking, hypertension, and sex on the phenotypic expression of familial intracranial aneurysms in siblings. *Neurosurgery* 2001; 48:64-9.
261. Hop JW, Rinkel GJ, Algra A, van Gijn J. Case-fatality rates and functional outcome after subarachnoid hemorrhage: a systematic review. *Stroke* 1997; 28:660-64.
262. Puddey IB, Beilin LJ, Vandongen R, Rouse IL, Rogers P. Evidence for a direct effect of alcohol consumption in normotensive men: a randomized controlled trial. *Hypertension* 1985;7:707-13.

- 
263. Intersalt Cooperative Research Group. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. *BMJ* 1988;297:319-28.
264. Ettinger PO, Wu CF, De La Cruz C Jr, Weisse AB, Ahmed SS, Regan TJ. Arrhythmias and the "Holiday Heart": alcohol-associated cardiac rhythm disorders. *Am Heart J* 1978;95:555-62.
265. Berglund M, Risberg J. Regional cerebral blood flow during alcohol withdrawal. *Arch Gen Psychiatry* 1981;38(3):351-5.
266. Klatskv AL, Friedman GD, Siegelaub AB, Gerard MJ. Alcohol consumption and blood pressure: Kaiser-Permanente multiphasic health examination data. *N Engl J Med* 1977;296:194-200.
267. Kataoka K, Taneda M, Asai T, Yamada Y. Difference in nature of ruptured and unruptured cerebral aneurysms. *Lancet* 2000;355:203.
268. Frösen J, Piippo A, Paetau A, Kangasniemi M, Niemela M, Hernesniemi J, et al. Remodelling of saccular cerebral artery aneurysm wall is associated with rupture: histological analysis of 24 unruptured and 42 ruptured cases. *Stroke* 2004;35:2287-93.
269. Pentimalli L, Modesti A, Vignati A, Marchese E, Albanese A, Di Rocco F, et al. Role of apoptosis in intracranial aneurysm rupture. *J Neurosurg* 2004;101:1018-25.
270. Tulamo R, Frösen J, Junnikkala S, Paetau A, Pitkäniemi J, Kangasniemi M, et al. Complement activation associates with saccular cerebral artery aneurysm wall degeneration and rupture. *Neurosurgery* 2006;59:1069-76.



271. Hashimoto T, Meng H, Young WL. Intracranial aneurysms: links among inflammation, hemodynamics and vascular remodeling. *Neurol Res* 2006;28:372-80.
272. Yasui N, Suzuki A, Nishimura H, Suzuki K, Abe T. Long-term follow-up study of unruptured intracranial aneurysms. *Neurosurgery* 1997;40:1155-59.
273. Wiebers DO, Whisnant JP, Sundt TM Jr, O'Fallon WM. The significance of unruptured intracranial saccular aneurysms. *J Neurosurg* 1987;66:23-9.
274. Sonobe M, Yamazaki T, Yonekura M, Kikuchi H. Small unruptured intracranial aneurysm verification study: SUAVe study, Japan. *Stroke* 2010;41:1969-77.
275. Allcock JM, Canham PB. Angiographic study of the growth of intracranial aneurysms. *J Neurosurg* 1976;45:617-21.
276. Burns JD, Huston J III, Layton KF, Piepgras DG, Brown RD Jr. Intracranial aneurysm enlargement on serial magnetic resonance angiography: frequency and risk factors. *Stroke* 2009;40:406-11.
277. Dispensa BP, Saloner DA, Acevedo-Bolton G, Achrol AS, Jou LD, McCulloch CE, et al. Estimation of fusiform intracranial aneurysm growth by serial magnetic resonance imaging. *J Magn Reson Imaging* 2007;26:177-83.
278. Juvela S. Natural history of unruptured intracranial aneurysms: risks for aneurysm formation, growth, and rupture. *Acta Neurochir Suppl* 2002; 82:27-30.
279. Koffijberg H, Buskens E, Algra A, Wermer MJ, Rinkel GJ. Growth rates of intracranial aneurysms: exploring constancy. *J Neurosurg* 109:176-185, 2008
280. Kamitani H, Masuzawa H, Kanazawa I, Kubo T. Bleeding risk in unruptured and residual cerebral aneurysms—angiographic annual growth rate in nineteen patients. *Acta Neurochir (Wien)* 1999;141:153-59.

281. Miyazawa N, Akiyama I, Yamagata Z. Risk factors for growth of unruptured intracranial aneurysms: follow-up study by serial 0.5-T magnetic resonance angiography. *Neurosurgery* 2006;58:1047–53.
282. Phan TG, Huston J III, Brown RD Jr, Wiebers DO, Piepgras DG. Intracranial saccular aneurysm enlargement determined using serial magnetic resonance angiography. *J Neurosurg* 2002;97:1023–28.
283. Inoue T, Shimizu H, Fujimura M, Saito A, Tominaga T. Annual rupture risk of growing unruptured cerebral aneurysms detected by magnetic resonance angiography. *J Neurosurg* 2012;117:20-5.
284. Kojima T, Waga S. More than one aneurysm on the same artery. *Surg Neurol* 1984;22:403–08.
285. Mckissock W, Richardson A, Walsh L, Owen E. Multiple intracranial aneurysms. *Lancet* 1964;1:623-26.
286. Rinne J, Hernesniemi J, Puranen M, Saari T. Multiple intracranial aneurysms in a defined population: prospective angiographic and clinical study. *Neurosurg* 1994;35:803-8.
287. Locksley HB, Sahs AL, Sandler R. Report on the cooperative study of intracranial aneurysms and subarachnoid hemorrhage. 3. Subarachnoid hemorrhage unrelated to intracranial aneurysm and A-V malformation. A study of associated diseases and prognosis. *J Neurosurg* 1966;24:1034-56.
288. Sato O, Kanazawa I, Kokunai T, Kobayashi M. Seven intracranial aneurysms of the internal carotid artery. Diagnosis by magnification angioautotomography. *Neuroradiol* 1978;15:189–92.

- 
289. Rinne J, Hernesniemi J, Niskanen M, Vapalahti M. Management outcome for multiple intracranial aneurysms. *Neurosurgery* 1995;36:31-7.
290. Juvela S. Risk factors for multiple intracranial aneurysms. *Stroke* 2000;31:392-97.
291. Wermer MJ, van der Schaaf IC, Algra A, Rinkel GJ. Risk of rupture of unruptured intracranial aneurysms in relation to patient and aneurysm characteristics: an updated meta-analysis. *Stroke* 2007;38:1404-10.
292. Morita A, Fujiwara S, Hashi K, Ohtsu H, Kirino T. Risk of rupture associated with intact cerebral aneurysms in the Japanese population: a systematic review of the literature from Japan. *J Neurosurg* 2005;102:601-6.
293. Raghavan ML, Ma B, Harbaugh RE. Quantified aneurysm shape and rupture risk. *J Neurosurg* 2005;102:355-62.
294. Ujiie H, Tachibana H, Hiramatsu O, Hazel AL, Matsumoto T, Ogasawara Y, et al. Effects of size and shape (aspect ratio) on the hemodynamics of saccular aneurysms: a possible index for surgical treatment of intracranial aneurysms. *Neurosurgery* 1999;45:119-29.
295. van Gijn J, Kerr RS, Rinkel GJ. Subarachnoid haemorrhage. *Lancet* 2007;369:306-18.
296. Nehls DG, Flom RA, Carter LP, Spetzler RF. Multiple intracranial aneurysms: determining the site of rupture. *J Neurosurg* 1985;63: 342-48.
297. UCAS Japan Investigators, Morita A, Kirino T, Hashi K, Aoki N, Fukuhara S, et al. The natural course of unruptured cerebral aneurysms in a Japanese cohort. *N Engl J Med* 2012;366:2474-82.

298. Akagawa H, Kasuya H, Onda H, Yoneyama T, Sasahara A, Kim CJ, et al. Influence of endothelial nitric oxide synthase T-786C single nucleotide polymorphism on aneurysm size. *J Neurosurg* 2005;102:68-71.
299. Wang H, Kuwata S, Juji T, Yanagisawa M, Tokunaga K, Horie S, et al. Ethnic differences in allele frequencies of two microsatellite markers closely linked to the locus for polycystic kidney disease 1 (PKD1). *Hum Hered* 1995b;45:84-9.
300. Inoue K, Asao T, Shimada T. Ethnic-related differences in the frequency distribution of genetic polymorphisms in the CYP1A1 and CYP1B1 genes in Japanese and Caucasian populations. *Xenobiotica* 2000; 30:285-95.
301. Ko NU, Rajendran P, Kim H, Rutkowski M, Pawlikowska L, Kwok PY, et al. Endothelial nitric oxide synthase polymorphism (-786T->C) and increased risk of angiographic vasospasm after aneurysmal subarachnoid hemorrhage. *Stroke* 2008;39:1103-08.
302. Pena SD, Bastos-Rodrigues L, Pimenta JR, Bydlowski SP. DNA tests probe the genomic ancestry of Brazilians. *Braz J Med Biol Res* 2009;42:870-6.
303. Khurana VG, Sohni YR, Mangrum WI, McClelland RL, O’Kane DJ, Meyer FB, et al. Endothelial nitric oxide synthase T-786C single nucleotide polymorphism: a putative genetic marker differentiating small versus large ruptured intracranial aneurysms. *Stroke* 2003;34: 2555–59.
304. Khurana VG, Sohni YR, Mangrum WI, McClelland RL, O’Kane DJ, Meyer FB, et al. Endothelial nitric oxide synthase gene polymorphisms predict susceptibility to aneurysmal subarachnoid haemorrhage and cerebral vasospasm. *J Cereb Blood Flow Metab* 2004b;24: 291–97.

305. Rudic RD, Shesely EG, Maeda N, Smithies O, Segal SS, Sessa WC. Direct evidence for the importance of endothelium-derived nitric oxide in vascular remodelling. *J Clin Invest* 1998;101: 731–36.
306. Paik DC, Ramey WG, Dillon J, Tilson MD. The nitrite=elastin reaction: implications for in vivo degenerative effects. *Connect Tissue Res* 1997;36: 241–51.
307. Johanning JM, Franklin DP, Han DC, Carey DJ, Elmore JR. Inhibition of inducible nitric oxide synthase limits nitric oxide production and experimental aneurysm expansion. *J Vasc Surg* 2001;33:579–86.
308. Tesauro M, Thompson WC, Rogliani P, Chaudhary PP, Moss J. Intracellular processing of endothelial nitric oxide synthase isoforms associated with differences in severity of cardiopulmonary diseases: cleavage of proteins with aspartate vs. glutamate at position 298. *Proc Natl Acad Sci USA* 2000;97: 2832–35.
309. Tsukada T, Yokoyama K, Arai T, Takemoto F, Hara S, Yamada A, et al. Evidence of association of the eNOS gene polymorphism with plasma NO metabolite levels in humans. *Biochem Biophys Res Com* 1998;245: 190–93.
310. Moon J, Yoon S, Kim E, Shin C, Jo SA, Jo I. Lack of evidence for contribution of Glu298Asp (G894T) polymorphism of endothelial nitric oxide synthase gene to plasma nitric oxide levels. *Thromb Res* 2002;107: 129–34.
311. Song MK, Kim MK, Kim TS, Joo SP, Park MS, Kim BC, et al. Endothelial nitric oxide gene T-786C polymorphism and subarachnoid hemorrhage in Korean population. *J Korean Med Sci* 2006; 21: 922-26.
312. Carmichael R. The pathogenesis of noninflammatory cerebral aneurysms. *J Pathol Bacteriol* 1950;62:1-19.
313. Stehbens WE. Histopathology of cerebral aneurysms. *Arch Neurol* 1963;8:272-85.

314. Folkman J, D'Amore PA. Blood vessel formation: what is its molecular basis? *Cell* 1996;87:1153-5.
315. Schwartz SM, deBlois D, O'Brien ER. The intima. Soil for atherosclerosis and restenosis. *Circ Res* 1995;77:445-65.
316. Owens GK. Regulation of differentiation of vascular smooth muscle cells. *Physiol Rev* 1995;75:487-517.
317. McAllister KA, Grogg KM, Johnson DW, Gallione CJ, Baldwin MA, Jackson CE, et al. Endoglin, a TGF-beta binding protein of endothelial cells, is the gene for hereditary haemorrhagic telangiectasia type 1. *Nat Genet* 1994;8(4):345-51.
318. Alberts MJ, Davis JP, Graffagnino C, McClenny C, DeLong D, Granger C, et al. Endoglin gene polymorphism as a risk factor for sporadic intracerebral hemorrhage. *Ann Neurol* 1997;41:683-86.
319. Peters DG, Kassam AB, Chang YF. A DNA sequence polymorphism in the endoglin gene is not associated with intracranial aneurysm or aneurismal subarachnoid hemorrhage. *Cerebrovasc Dis* 2005; 20:96-100.
320. Leikina E, Merts MV, Kuznetsova N, Leikin S. Type I collagen is thermally unstable at body temperature. *Proc Natl Acad Sci U S A* 2002;99:1314-8.
321. Servos S, Zachary I, Martin JF. VEGF modulates NO production: the basis of a cytoprotective effect? *Cardiovasc Res* 1999;41:509-10.
322. Wu X, Li D, Liu Z, Wan X, Wu Y, Jiang C, Qian Q. Vascular endothelial growth factor 1498C/T, 936C/T polymorphisms associated with increased risk of colorectal adenoma: a Chinese case-control study. *Mol Biol Rep* 2011;38:1949-55.

323. Frösen J, Piippo A, Paetau A, Kangasniemi M, Niemelä M, Hernesniemi J, et al. Growth factor receptor expression and remodeling of saccular cerebral artery aneurysm walls: implications for biological therapy preventing rupture. *Neurosurgery* 2006;58:534-41; discussion 534-41.
324. Grosskreutz CL, Anand-Apte B, Dupláa C, Quinn TP, Terman BI, Zetter B, D'Amore PA. Vascular endothelial growth factor-induced migration of vascular smooth muscle cells in vitro. *Microvasc Res* 1999;58:128-36.
325. Wittke-Thompson JK, Pluzhnikov A, Cox NJ. Rational inferences about departures from Hardy-Weinberg equilibrium. *Am J Hum Genet.* 2005;76:967-86.
326. De Rooij NK, Linn FH, van der Plas JA, Algra A, Rinkel GJ. Incidence of subarachnoid haemorrhage: a systematic review with emphasis on region, age, gender and time trends. *J Neurol Neurosurg Psychiatry* 2007;78:1365-72.
327. Longstreth WT Jr, Nelson LM, Koepsell TD, van Belle G. Cigarette smoking, alcohol use, and subarachnoid hemorrhage. *Stroke* 1992;23:1242-49.
328. Ruigrok YM, Rinkel GJ, Wijmenga C. Genetics of intracranial aneurysms. *Lancet Neurol* 2005; 4:179–89.
329. Isaksen J, Egge A, Waterloo K, Romner B, Ingebrigtsen T. Risk factors for aneurysmal subarachnoid haemorrhage: the Tromsø study. *J Neurol Neurosurg Psychiatry* 2002;73:185-7.
330. Anderson C, Ni Mhurchu C, Scott D, Bennett D, Jamrozik K, Hankey G Australasian Cooperative Research on Subarachnoid Hemorrhage Study Group. Triggers of subarachnoid hemorrhage: role of physical exertion, smoking, and alcohol in the Australasian Cooperative Research on Subarachnoid Hemorrhage Study (ACROSS). *Stroke* 2003;34:1771-76.

- 
331. Juvela S. Prevalence of risk factors in spontaneous intracerebral hemorrhage and aneurismal subarachnoid hemorrhage. *Arch Neurol* 1996;53:734-40.
  332. Ruigrok YM, Buskens E, Rinkel GJE. Attributable risk of common and rare determinants of subarachnoid hemorrhage. *Stroke* 2001; 32: 1173–75.